

# Cardiac Strain and Left Ventricular Mass Quantification for Hypertrophic Cardiomyopathy using 4D Echocardiography

Tam, Hannah

**Background:** Strain and Left ventricular mass (LVM) determination are essential for the evaluation of the progress and function of hypertrophic cardiomyopathy (HCM). Free of geometric assumptions and angle dependencies, four-dimensional echocardiography (4DE) has been shown to be a superior method in determining myocardial strain and LVM than current conventional methods. However, its feasibility in determining myocardial strain and LVM for HCM has not been reported.

**Methods:** A balloon was sutured into the left ventricle (LV) of 18 porcine hearts and connected to a calibrated pump. The right ventricle free wall was sectioned into quarters and sutured on the septum one at a time to simulate septal wall thickness. 4D and Motion-mode images were obtained at baseline and each simulated HCM level. Full-volume 4D loops were analyzed to quantify global longitudinal, circumferential, radial, and area strain, as well as the regional strain values at the hypertrophied mid-septum segment. **Results:** Full-volume 4D loops were analyzed; in HCM models, 4D-derived LVM demonstrated superior linear correlation with displacement values ( $R^2=0.95$ ) when compared to M-mode ( $R^2=0.55$ ). However, both methods demonstrated excellent correlations with displacement values (both  $R^2>0.90$ ,  $p<0.001$ ) at baseline. Bland-Altman analyses showed an overestimation in both methods. All global strain and regional strain (MS) were significantly reduced in HCM models when compared to the baseline (all  $p$ -values  $<0.05$ ), signifying decreased function and thereby differentiating physiological and pathological hypertrophy. **Conclusions:** Full-volume 4DE is a feasible method for left ventricular mass detection and global and regional strain determination in HCM with abnormal septal wall.