

Counteracting Pb²⁺ Induced Neurodegeneration: A Chemical and Genetic Approach towards Enhanced Synaptic Plasticity

Fiacco, Jason

Sadhu, Kevin

The neurotrophin BDNF has been shown to engage in retrograde signaling and interneuronal transport, causing increased synaptic transmission and neurogenesis, respectively. It was hypothesized that Pb²⁺ exposure causes neurodegeneration by lowering BDNF levels and transport. Thus, improving BDNF levels and transport would counteract Pb²⁺ induced neurodegeneration. To increase BDNF levels, Pb²⁺ exposed *Drosophila* were treated with curcumin, and dissected to isolate their neuromuscular junctions. Fluorescent indicators FRET, FM 1-43 dye, and FURA-2 dye were used to analyze levels of cAMP, presynaptic vesicle rate, and Ca²⁺ influx respectively. To increase neurotrophin transport, a knockout of FOS-1 was induced in Pb²⁺ exposed *C. elegans* with GFP-tagged motor neurons, and were then observed with a Zeiss-Axiovert fluorescent microscope and analyzed using ImageJ to quantify neurogenesis. Locomotion and lifespan were also analyzed in both flies and worms. The results suggest that curcumin caused an increase in Ca²⁺ influx, vesicle rate, and locomotion, but not in cAMP or lifespan. Knockout of FOS-1 increased neuronal growth (GFP) and lifespan, but not locomotion. This suggested that improving BDNF levels and transport was effective in counteracting neurodegeneration, thus indicating that Pb²⁺ causes neurodegeneration largely through its detrimental effects on BDNF. Furthermore, a third pathway involving acetylcholine was impaired, suggesting additional consequences of Pb²⁺ exposure. It was therefore determined that the inhibition of BDNF by Pb²⁺ significantly deteriorates motor functions, synaptic functions and lifespan, proposing this pathway as a suitable target when counteracting Pb²⁺ neurodegeneration.

Awards Won:

Second Award of \$2,000