

Biosorption of Manganese Mine Effluents Using Crude Chitin from Shell Wastes of Philippine Bivalves

Molejona Jr., Randy (School: Iloilo National High School)

Saquin, Elaine Nicole (School: Iloilo National High School)

The area around Ajuy river in Iloilo, Philippines is currently being mined for manganese ore, and river water samples exceed the maximum manganese contaminant level set by US-EPA. At the same time, the surplus of local bivalve waste is another environmental concern. Studies show that chemical treatment compromises water quality leaving toxic residues, and an alternative treatment process is biosorption, or using the physical and chemical properties of a biomass to adsorb heavy metals in contaminated water. The study aims to extract crude chitin from shell wastes of *Bractechlamys vexillum*, *Perna viridis*, and *Placuna placenta* and determine its adsorption capacity on manganese in simulated and actual mine water. Crude chitin was obtained by pulverization, deproteinization, demineralization, and decolorization of shells. Biosorption by flocculation followed 5 g: 50 mL chitin-to-water ratio. Filtrates were analyzed using MP-AES after 24 hours. In both actual and simulated mine water respectively, *B. vexillum* yielded the highest adsorption percentage of 91.43% and 99.58%, comparable to *P. placenta* of 91.43% and 99.37%, while significantly different to *P. viridis* of -57.14% and 31.53%, ($p < 0.05$). FT-IR validated the presence of chitin in shells based on carbonyl-containing functional groups at peaks 1530-1560 cm^{-1} and 1660-1680 cm^{-1} . SEM micrographs showed the amorphous and non-homogenous structure of chitin. Thus, crude chitin from *B. vexillum* and *P. placenta* can be bio-sorbents for water treatment of manganese-impacted effluents, and promote appropriate waste management of local bivalves.

Awards Won:

Third Award of \$1,000