

# On a Variation of the Witsenhausen Problem Concerning Maximal $\pi/2$ -Avoiding Spherical Sets

Perez, AnaMaria (School: Albuquerque Academy)

The problem of finding the largest  $\pi/2$ -avoiding spherical set first appeared in the AMS monthly journal in 1974. H. S. Witsenhausen asked readers to determine  $\alpha(n) = (|U|)/(|S^{n-1}|)$ , where  $U$  is the largest  $\pi/2$ -avoiding set in the  $n$  dimensional sphere  $S^{n-1}$ , that is, the largest set on the sphere in  $n$  dimensions containing no orthogonal vectors. We look at a variation of this problem: determine  $\alpha(n,k) = (|U|)/(|S^{n-1}|)$ , where  $U$  is the largest set on the  $n$  dimensional sphere  $S^{n-1}$  that contains no  $k$  mutually orthogonal vectors. We begin by specializing the problem to the case  $n=k=3$  by considering spherical sets that avoid three mutually orthogonal vectors in 3D in an attempt to determine  $\alpha(3,3)$  which allows for more interesting configurations and visualization. We find the largest spherical subset  $U$  not containing three mutually orthogonal vectors in the following cases: (i)  $U$  is a large double cap; (ii)  $U$  is a centered band; or (iii)  $U$  is a wedge shape. We then prove a generalized lower bound for  $\alpha(n,k)$  for any  $n \geq 3$ . By taking  $U$ , the union of two double caps and a centered band, we increase the area further. The measures of these sets give a provable lower bound for  $\alpha(3,3)$ . Finally, by smoothing the band using computational results, we find a configuration created by the intersection of elliptical cylinders with the sphere that gives an even better lower bound for  $\alpha(3,3)$ . We conjecture this to be the largest spherical set for  $\alpha(3,3)$ , and one that can be generalized to higher dimensions.