

Combating Antibiotic-Resistant Bacteria With Leptospermum Honey-Incorporated Calcium Alginate Dressing

Chung, Ting Yan (School: St. Paul's Convent School)

Liu, Tsz Ki (School: St. Paul's Convent School)

Over the past several decades, excessive antibiotic use has led to the rapid development of antibiotic-resistant bacteria. The decreasing effectiveness of antibiotics in managing chronic wounds has quickened, thus threatening patient health. Given honey's long history in wound treatment, this study had two goals: to assess the antimicrobial activity of honey; and develop an absorbent, biodegradable and antibacterial honey-based wound dressing. We compared the antimicrobial effect of honey from different floral sources and investigated the dose-dependent effect of the *Leptospermum* honey using agar well diffusion assays. In this procedure, the agar plate surface was inoculated with skin surface microorganisms, samples were introduced into wells, and plates were incubated at 28°C for 24 hours. Zones of inhibition were then calculated using NIH ImageJ. Among the 6 different floral sources tested, honey derived from *Leptospermum scoparium* exhibited the most potent antimicrobial ability. There was a positive correlation ($r=0.972$, $p=0.006$) between the concentration of *Leptospermum* honey and inhibited microbial growth. The hydrogel product is created by, firstly, integrating honey into sodium alginate, followed by a reaction with calcium chloride, which traps honey within the net structure of the polymer. From the swelling capacity experiment, honey-based hydrogel showed a maximum uptake of 25% of its original mass. From the biodegradability test, honey-based hydrogel demonstrated a biodegradation level >14% greater than the control hydrocolloid within the first week. This dressing is antibacterial, absorbent and biodegradable, and has the potential to substitute topical wound agents.