

Research on the Microscopic Mechanism of CO₂ Capture Using TBAB Semi-Clathrate Hydrate Formation

Lu, Guoxiang (School: Chongqing No 1 International Studies School)

Purpose of my research (1) Use a new additive to form semi-clathrate hydrate at lower pressures and higher temperatures for CO₂ capture. (Research question) (2) Can Tetra-n-butyl Ammonium Bromide (TBAB) work? (3) Understand the mechanism of CO₂ capture by TBAB semi-clathrate from different aspects (microscopic morphology, Raman spectroscopy, etc.) (4) Find out if TBAB semi-clathrate can capture CO₂ efficiently

Experimental procedure High-pressure DSC Measure the temperature and pressure data for CO₂ hydrate formation. Find out the thermodynamic conditions for TBAB + CO₂ semi-clathrate hydrate formation. Microscopic experiment Explore the morphology of TBAB + CO₂ semi-clathrate by recording the microscopic images. Obtain the microscopic images of TBAB+CO₂ semi-clathrate hydrate. Understand the hydrate growth behavior. Raman experiment Measure Raman spectral intensity and determine the structure transition of TBAB + CO₂ semi-clathrate hydrate. Disclose the mechanism of CO₂ capture on the molecular scale.

Conclusions It is found that TBAB is an effective additive which can form semi-clathrate hydrate for CO₂ capture at lower pressures and higher temperatures. The microscopic observation and Raman spectroscopy experiments indicate that 2.57 mol% TBAB is the optimal concentration for CO₂ capture. The fastest rate of TBAB+CO₂ hydrate is obtained at 2.57 mol% TBAB, and the largest amount of CO₂ is captured at this TBAB concentration

Awards Won:

YM American Academy: Third Award of \$500.00