

Computational Modeling of CO₂ Migration and Trapping in Geologic Formations for Risk Assessment of Carbon Sequestration Impacting Carbon Footprint

Pandey, Shaunak (School: Vandegrift High School)

Over the last 150 years, the average global temperature on Earth has rapidly increased. This detrimental impact has created—among others—widespread famine, the melting of ice caps, larger, more frequent storms, drought, and poverty. Geologic carbon sequestration (GCS) involves the process of artificially injecting CO₂ into underground geologic formations such as saline aquifers, enabling us to continue using fossil fuels as a source of energy while mitigating their environmental impacts, as it is unlikely that we will be able to transition to renewable energy sources overnight. This experiment leveraged computational fluid dynamics and high-performance computing to explore the mechanics of CO₂ transport and trapping in rough-walled fractures focusing on the effect of fracture geometric factors such as aperture, roughness, and wettability—aimed towards GCS leakage risk assessment to mitigate potential leakage into urban infrastructure for large-scale implementation. When the fracture aperture was tight, CO₂ bypassed the water in the high capillary pressure zone. While for a wide water-wet fracture, CO₂ tends to centrally migrate to the fracture opening and thin film of residually trapped water occurs on the surface of the fracture. Less water-wet conditions led to a larger degree of CO₂ capillary trapping on the surface of the fracture. An unstable and fingering type of CO₂ displacement by increasing the injection rate in a rough fracture indicates that the fracture roughness is a key factor in controlling the CO₂ displacement regime in potentially fractured reservoirs. The results of this study provide novel insight of CO₂ behavior in rough-walled fractures.

Awards Won:

University of Texas at Dallas: Back-up scholarship recipients