

Management of Nitrogen and Phosphorus Biogeochemical Cycles Through Harvest of *Chlorella vulgaris* and *Elodea canadensis*

Kemp, Malory (School: Cavalier Public High School)

Biogeochemical cycles are series of processes that transfer compounds through an ecosystem. Can the geochemical cycles of phosphorus and nitrogen be managed effectively through harvest of *Chlorella Vulgaris* and *Elodea Canadensis*? Can the harvested bio-materials be used to recycle nitrogen and phosphorus into the growing cycle of crops such as rye? Can *Chlorella Vulgaris* algal balls be utilized for easier capture and application of nutrient rich algae? Can nitrogen and phosphorus used by *Elodea Canadensis* be recycled as a fertilizer? I hypothesize that *Chlorella Vulgaris* and *Elodea Canadensis* will be able to effectively lower nitrogen and phosphorus levels in closed pond systems. Further, I hypothesis that *Chlorella Vulgaris* and *Elodea Canadensis* can be incorporated into soils as fertilizer for use in grain crops such as rye. The biomass of the harvested rye grass after 10 and 17 days ranged from an average of 9.23/5.40 grams for *Elodea Canadensis*, 11.86/6.28 grams for *Chlorella Vulgaris*, 9.83/6.59 *Chlorella Vulgaris* algal balls, and Control average of 6.12/3.90 grams. Total biomass harvest ranged from an average of 14.63 grams for *Elodea Canadensis*, 18.14 grams for *Chlorella Vulgaris*, 16.42 *Chlorella Vulgaris* algal balls, and Control average of 10.02 grams. Results can not adequately support that *Chlorella Vulgaris* and *Elodea Canadensis* will be able to effectively lower nitrogen and phosphorus levels in closed pond systems. Further study over a longer period of time may track the levels of nutrients and the effectiveness of *Chlorella Vulgaris* and *Elodea Canadensis*. I accept my hypothesis that *Chlorella Vulgaris* and *Elodea Canadensis* can be incorporated as fertilizer for use in grain crops such as rye.