

# The Group Commutator Length in Terms of Group Ring

Vagner, Maksim

Solikov, Pavel

Let  $G$  be a group and  $[G, G]$  be its commutator subgroup which is generated by special elements named commutators. Therefore, every element from the commutator subgroup can be represented as a product of commutators. Such representation of an element  $g$  from  $[G, G]$  that contains the least possible number of commutators is called a minimal commutator representation of  $g$ . The number of commutators in a minimal representation of  $g$  is called the commutator length of  $g$ . The purpose of our work is creation of the instrument for transfer the work with commutator length from the group  $G$  to its group ring  $ZG$ . We give an equivalent definition of commutator length for element  $g$  from  $[G, G]$ , which does not use concept of group commutator, but uses concept of ring commutator in the group ring. Now we introduce the following concept. Suppose  $x, y$  is from  $ZG$ ; then the element  $xy - yx$  is called the ring commutator of  $x, y$ . Suppose  $g$  is from  $[G, G]$ ; then there exist a representation of element  $g - 1$  as sum of ring commutators, multiplied on elements from  $G$ . The least number of summands in the representation is called ring commutator length of element  $g - 1$ . The aim of this paper is to prove the equality commutator length of  $g$  and ring commutator length of  $g - 1$ . Therefore, ring commutator length is equal to the group commutator length. Consequently, it is another definition of the well-known object. We believe that the general commutator length theory will greatly benefit from using this definition.