

The Important Discovery of a Greenhouse Gas "Bio-Catcher": Carbon Fixation in Trentepohlia

Lu, Yiyang

Blood red boulders are scattered across a large area above the creek in Hailuoguo, Gongga Mountain, Sichuan, the red matter attached to the stones is a species of Trentepohlia, a type of subaerial filamentous green chlorophyte algae which can be used as a greenhouse gas "bio-catcher" in the future. In order to test the anti-stress ability and carbon fixing ability of Trentepohlia, chlorophyll fluorescence analysis and far-infrared gas analysis were used and I found that the algae's maximum photochemical productivity reached 0.67 ± 0.015 , and displayed no significant variation across a range of radiation intensities from 0 to $400 \mu\text{W cm}^{-2}$; it also possesses strong tolerance for UV radiation. At 10°C , with a light intensity of $45 \mu\text{mol m}^{-2} \text{s}^{-1}$, its carbon fixation capacity is $351.57 \mu\text{mol CO}_2 \text{mg}^{-1} \text{Chl a h}^{-1}$ —2.6 times that of Nostoc, and 1.4 times that of Chlorella under similar conditions, reflecting the algae's relatively strong capacity for fixing carbon dioxide. The algae's light compensation point of photosynthesis is $28 \mu\text{mol m}^{-2} \text{s}^{-1}$, and its light saturation point is $565 \mu\text{mol m}^{-2} \text{s}^{-1}$, light of intensity over $800 \mu\text{mol m}^{-2} \text{s}^{-1}$ triggers a sharp decline in carbon fixation, that means the cells have a light protection mechanism; Calculations using the data from testing reveal that the Trentepohlia algae in Hailuoguo sequester approximately 10 tons of CO_2 a year.