## The Optimization of Degradation of Azo Dyes via Saccharomyces cerevisiae

Segaran, Nicole (School: Carmel High School)

Azo dyes are commonly used in the textile industry, and are a major constituent of textile factory waste. However, azo-linkages remain non biodegradable, and are highly toxic, linked the neurosensory damage in aquatic biota and cancer. While previous studies show various yeasts and bacteria are able to cleave azo-dye linkages, several have been found to have negative effects on wildlife, and are difficult or costly to culture. Saccharomyces cerevisiae, is a possible biosorbent of dyes, and has potential to degrade azo dyes. This study contains three purposes: 1) confirm Saccharomyces Cerevisiae as a possible decolorizing agent of azo dyes 2) experiment with the Active Dry Yeast form of S. Cerevisiae as a degrading agent, in comparison to the wild strain, and 3) observe the mechanisms underlying degradation via S. Cerevisiae. Experimentation confirmed S. Cerevisiae as a possible degrading agent of azo dyes, decolorizing the dyes on average >70% (analyzed using spectrophotometry). Furthermore Active Dry Yeast was proven a more efficient degrading agent than the S. Cerevisiae used to degrade Reactive Red 2 was observed to be both bioaccumulation and biosorption due to the characteristics analyzed during experimentation; however, further studies delving into the cellular enzymatic activity is recommended to confirm that both bioaccumulation and biosorption are present.