

Alpha-synuclein Enhances Toxicity of Tau Oligomers in vitro

Berland, Katelynne (School: New Horizons Governor's School for Science and Technology)

Aggregation of protein tau into neurofibrillary tangles and α -synuclein into Lewy bodies are hallmarks of Alzheimer's and Parkinson's disease, respectively. These proteins are known to form pre-fibrillar aggregates called oligomers (multiples of monomeric protein), which are considered the toxic entity of the diseases. The coexistence of both tau and α -synuclein is associated with worse disease progression, which suggests a synergistic relationship between these proteins. Although the interaction between α -synuclein and tau oligomers has been demonstrated in Parkinson's and Dementia with Lewy body (DLB) cases, the mechanism of toxicity remains unclear. To probe this synergistic mechanism, a stable cell line was created with expression of full length human α -synuclein in SH-SY5Y neuroblastoma cells. siRNA was used to knockdown the expression of α -synuclein in SH-SY5Y cells prior to toxicity assay. The stable cell line and SH-SY5Y cells were exposed to recombinant oligomeric species of tau or vehicle (PBS) for 8-24 hours. Immunostaining and MTT assay were used to assess cell viability. RT-PCR was used to assess mRNA expression following the toxicity assay. Results depict that the knockdown of α -synuclein prevents tau oligomer toxicity in vitro. Furthermore, the coexistence of both proteins enhances the neurotoxicity of the tau oligomers. These findings strongly suggest that a synergistic mechanism among tau and α -synuclein potentiates toxicity. Elucidating the mechanisms of neurodegeneration could reveal novel protein targets of disease progression.

Awards Won:

Fourth Award of \$500