

Galleria mellonella as a Proxy for the Human Immune Response: Allium sativum's Effectiveness Against Enterobacter cloacae

Chang, Leo (School: Central Virginia Governor's School for Science and Technology)

This study aimed to evaluate the effectiveness of garlic extract against *E. cloacae* infections. First, inhibition zones were created using lawns of *E. cloacae* and sterile disks soaked with decreasing concentrations of garlic extract. A one-way ANOVA ($\alpha = .05$, $p = 4.36 \times 10^{-27}$) followed by a Tukey test identified 100% garlic extract as the MIC. Seven trials involved *Galleria mellonella*, which has a comparable innate immune response to that of humans, being divided into five groups: *E. cloacae* control, PBS control, garlic extract control, prophylactic treatment, and therapeutic treatment. Larvae were injected with 10 μ l of inocula, with bacteria and garlic extract injections separated by two hours for treatment groups. Survival was measured every 24 hours over 72 hours. A one-way ANOVA ($p = 2.41 \times 10^{-13}$) and Tukey test revealed significant differences between therapeutic and prophylactic treatments. A two-sample t-test between prophylactic and therapeutic treatment groups ($p = .007$) confirmed the higher effectiveness of therapeutic treatment. A parallel experiment collected hemolymph from larvae every 24 hours after the last injection starting at the 2-hour mark. A one-way ANOVA ($p = 6.31 \times 10^{-31}$) and post-hoc test highlighted significant differences between treatments, supported by a two-sample t-test ($p = 3.74 \times 10^{-8}$), again confirming the superior efficacy of therapeutic treatment. These findings validated the H1 hypothesis but did not support the H2 and H3 hypotheses which suggested the prophylactic treatment would increase survival rate and decrease PO in hemolymph. The results suggest that while prophylactic treatment may not be effective, therapeutic treatment should be researched further.

Awards Won:

